

Concept paper

Refining the process of agent selection through understanding plant demography and plant response to herbivory

S Raghu,^{1*} John R Wilson² and K Dhileepan¹

¹Cooperative Research Centre for Australian Weed Management, Alan Fletcher Research Station, Queensland Department of Natural Resources and Mines, PO Box 36, Sherwood, Qld 4075, Australia.

²Centre for Invasion Biology, University of Stellenbosch, Private Bag X1, Matieland 7600, South Africa.

Abstract

Understanding plant demography and plant response to herbivory is critical to the selection of effective weed biological control agents. We adopt the metaphor of ‘filters’ to suggest how agent prioritisation may be improved to narrow our choices down to those likely to be most effective in achieving the desired weed management outcome. Models can serve to capture our level of knowledge (or ignorance) about our study system and we illustrate how one type of modelling approach (matrix models) may be useful in identifying the weak link in a plant life cycle by using a hypothetical and an actual weed example (*Parkinsonia aculeata*). Once the vulnerable stage has been identified we propose that studying plant response to herbivory (simulated and/or actual) can help identify the guilds of herbivores to which a plant is most likely to succumb. Taking only potentially effective agents through the filter of host specificity may improve the chances of releasing safe and effective agents. The methods we outline may not always lead us definitively to the successful agent(s), but such an empirical, data-driven approach will make the basis for agent selection explicit and serve as testable hypotheses once agents are released.

Key words agent efficacy, matrix models, *Parkinsonia aculeata*, plant demography, resistance, susceptibility.

INTRODUCTION

Invasive species pose significant threats to biodiversity and primary production (Mack *et al.* 2000; Pimentel *et al.* 2001; Pimentel 2002). Hence regulatory authorities and society in general is becoming risk-averse to introduction of any organisms into a region where they do not naturally occur. This has significant implications for the practice of classical biological control that relies on the introduction of specialist insects and pathogens into novel environments to manage populations of invasive species (Howarth 1991; McFadyen 1998; Sheppard *et al.* 2003; Sheppard & Raghu 2005). There is increasing demand for better evaluation of the risks and benefits of classical biological control agents prior to their introduction (Howarth 1991; Simberloff & Stiling 1996; McEvoy & Coombs 2000; Louda *et al.* 2003a,b; Sheppard 2003; Louda & Stiling 2004; McClay & Balciunas 2006).

Although it is widely acknowledged that classical biological control can be a valuable tool in the management of invasive plants (McFadyen 1998; Thomas & Willis 1998; McEvoy 2002; Hoddle 2004a,b), there is a risk (and an economic cost) attached to introducing a non-native organism. These risks and

costs can be reduced if we devise methods that limit the release of ineffective agents. Rather than revisit debates about whether we can be more efficient/rigorous about the way we make agent selection decisions in biological control (e.g. Schroeder & Goeden 1986; Cullen 1992, 1995; Denoth *et al.* 2002; McFadyen 2003; Sheppard 2003), we outline methods that we think could improve the chances of releasing effective agents (Fig. 1). We propose that subjecting the pool of potential agents identified in native range surveys to the filters of plant demography (Briese & Zapater 2002; Briese *et al.* 2002a; Sheppard 2003; Briese 2004; Buckley *et al.* 2005), and responses to herbivory (Broughton 2003; van Kleunen *et al.* 2004; Raghu & Dhileepan 2005) can significantly enhance the likelihood of selecting effective agents. These agents can subsequently be taken through the filter of host-specificity testing (Fig. 1).

PLANT DEMOGRAPHY (FILTER 1)

The structure of plant populations (genetic, spatial, age, size, stage structure) can be used to effectively direct agent selection (Silvertown & Lovett Doust 1993; Crawley 1997). Our objective in weed biological control is to use host-specific insects or pathogens to reduce the growth and/or spread of weed populations to achieve our *a priori* definition of

*Present address: Centre for Ecological Entomology, Illinois Natural History Survey & University of Illinois, 1816, S. Oak St., Champaign, IL 61820, USA (email: raghu@uiuc.edu).

‘successful management’ (van Klinken & Raghu 2006). Ideally we would identify the critical stages that influence the target weed’s demography by comparing detailed plant demographic data between the native and invaded range (McEvoy & Coombs 1999; Buckley *et al.* 2005). However, such data are not always available, or logistically possible to collect, in the early stages of a biological control program when agent selection decisions are made. In such circumstances, developing relatively simple models of the target weed’s life history based on sound natural history information and any available data on the target (or closely related species) may be extremely valuable.

Matrix models as tools in identifying the Achilles heel of weed populations

Models have been used as tools to understand plant herbivore interactions. In particular matrix models (Caswell 2001) have been used extensively in ecology to identify the most important factors determining a species’ population growth or spread. In the context of weed management such models can help identify the vulnerable stage in the weed’s life cycle.





Fig. 1. Nested filters in the agent selection process based on plant (weed) ecology.

Fig. 2. A hypothetical life cycle of a tree. The numbers adjacent to the arrows are the probabilities of an individual moving from one stage to the next. Because individuals survive, grow or die, the probabilities from each stage add up to 1 (except for reproducing individuals where one tree produces many seeds, for example, 10 in this case).



The first step in building a matrix population model is to split the population into a series of non-overlapping classes (e.g. age, size or stage classes) that captures the population structure of the particular weed. Then you need to measure how individuals move between stages (Fig. 2). These parameters can be estimated from tagging and following individuals at a site from year to year (or from month to month for annual species). Alternatively, this may be approximated based on expert opinion and sound natural history observations. The obvious caveat is that the inferences made from the model are only as reliable as the quality of the data and assumptions that went into its construction (e.g. how well the stages represent reasonable ‘thresholds’, and how the probabilities are fixed with time, space, competitive environment, etc.) (Caswell 2001). A simple example is shown in Figure 2, and the probabilities of moving between stages can be written in matrix form (a transition matrix, Fig. 3).

The population growth rate tends towards a constant and can be determined from the matrix by the dominant eigenvalue. Sensitivity (varying matrix elements by a fixed amount) and elasticity (varying matrix elements by a fixed proportion) analyses indicate the importance of each parameter to the population growth rate (Caswell 2001). Using these analyses we can assess the impact of various control methods (including biological control) on the weed. For example, the thistle *Carduus nutans* was predicted not to be controlled at one site in New Zealand by seed-feeding biological control agents unless additional measures, targeting other life stages, were in place (Shea & Kelly 1998). Equally, such a modelling approach can also be used to assess the impacts on non-target

	Seeds	Seedlings	Saplings	Mature trees
Seeds	0.8	0	0	10
Seedlings	0.1	0.2	0	0
Saplings	0	0.1	0.7	0
Mature trees	0	0	0.2	0.8

Fig. 3. A transition matrix for the hypothetical tree in Figure 2. The elements are colour-coded to match Figure 2 (red indicates survival rates, blue indicates development rates, green indicates reproduction).

plants by an insect under consideration as a biological control agent (Louda *et al.* 2005).

Constructing a matrix model of *Parkinsonia aculeata*

As an example of using matrix population models in weed management, we present a model developed for *Parkinsonia aculeata*, a rangeland weed of national significance in Australia.

Identifying discrete life-history stages. The first decision is how to split the population into discrete stages. Obviously seeds fall into a convenient category. However, it is less clear how to separate seedlings and adult plants – by age, by height or by stem/trunk diameter? We chose height as it is a good predictor of seed production and a tree's ability to survive frost and fire. Seedlings and adult plants were split into five categories: 0–1 m; 1–2 m; 2–3 m; 3–5 m; and greater than 5 m.

Calculating demographic parameters. Data collected from Alcoota Station in Central Australia in 2001/2 (van Klinken 2005; L Anderson and S Campbell unpubl. data 2005) were used to derive the different demographic parameters in the transition matrix.

Reproduction – Trees need to be taller than 1 m before they reproduce (Fig. 4a), and there is a clear increase in seed production with height (Fig. 4b). The change in seed production with height is well described by a step function (Fig. 4b). The seed production of each of the height classes was estimated from these relationships (Fig. 5).

Development rate – About 8% of seeds are expected to germinate in any year. The probability of moving between stages (development rates) was estimated from the growth rate curve (Figs 4c, 5).

Survival – Smaller trees are less likely to survive frost (Fig. 4d) and so in the transition matrix (Fig. 5), annual mortality declines with plant size. The rate of decay of the seed bank was estimated at about 0.98 per year; this figure includes those seeds germinating at a rate of 0.08 per year (van Klinken 2005; L Anderson and S Campbell unpubl. data 2005).

Based on this transition matrix, the population is expected to grow at this site by 36% every year. The elasticity of this transition matrix can then be calculated (Fig. 6).

The elasticity analysis suggests that the weed is best controlled by removing large trees (>5 m) as they have a massive effect on the population growth rate (Fig. 6). Seed production would have to be reduced by a very high level to create the same effect as cutting down a few large trees. However, the population is sensitive to changes in seed germination rate and the rate of development of young trees.

Logical agent selection decisions can be made based on this information. It is unlikely that an agent is going to inflict mortality on trees greater than 5 m in height on its own. Perhaps an alternate method of control (mechanical/chemical) is required to remove this size class. The elasticity analysis also suggests that seed feeders in this system may not have an

	Seeds	P(0–1)	P(1–2)	P(2–3)	P(3–5)	P(>5)
Seeds	0.02	0	2	86	1206	7586
P(0–1)	0.08	0.02	0	0	0	0
P(1–2)	0	0.08	0.105	0	0	0
P(2–3)	0	0	0.245	0.15	0	0
P(3–5)	0	0	0	0.35	0.56	0
P(>5)	0	0	0	0	0.14	0.95

Fig. 5. Transition matrix for a *Parkinsonia aculeata* population. The elements are colour-coded (red indicates survival rates, blue indicates development rates, green indicates reproduction).




Fig. 4. Effect of tree height on (a) onset of flowering; (b) seed production; (c) growth rate; (d) survival rate. Reproduction is well described by a step function, where the smooth line is the mean and the dotted line is the 95% confidence interval (CI). The other lines are fitted lines, which are used to estimate the parameters of the matrix model for *Parkinsonia aculeata*.

important effect on plant population dynamics. However, the above analysis suggests that affecting the development of young trees can significantly affect population growth rate. This information could serve as the *a priori* basis for selecting biological control agents.

This exercise illustrates how easily a simple model can be used to make predictions in weed management. However, it is important to remember that models are abstractions of the real world. Population dynamics differ year to year and site to site (Shea *et al.* 2005), as is certainly the case for *P. aculeata* in Australia (R van Klinken unpubl. data 2005). Ideally we would like to be able to construct mechanistic models for every system we work on (and there are some excellent programs that assist this process (e.g. DYMEX (Maywald *et al.* 1999)), but in most cases the data requirements are prohibitive. Therefore, matrix models still have a lot to offer as a first step.

The basic matrix models have been adapted in many ways by adding stochastic simulations using a probability distribu-

tion for each transition, calculating economic costs so elasticity analysis looks at how spending more money will affect population growth rate, including a matrix of dispersal to look at the spread of invasive species, and including density dependence so that the populations show sensible regulation (Caswell 2001). The main advantage of these models lies in their relative simplicity. They can be rigorously analysed without necessarily requiring large and precise data sets. This sort of approach lends itself quite amenable to the methods required in the process of agent selection.

Once the Achilles heel of the plant's life history has been identified, it is then possible to identify the type of damage that this weak link/transition is most likely to succumb to, by studying plant response to herbivory.

PLANT RESPONSE TO HERBIVORY (FILTER 2)

Plants experience damage from a wide variety of environmental factors (including herbivores), and exhibit a diversity of responses that can be broadly classified into susceptibility and resistance (Box 1; Belsky *et al.* 1993). From the perspective of selecting a biological control agent, we need to identify the types of damage (that a specialist herbivore can inflict) that results in the management of weed populations.

Ideally, we would like a biological control agent to elicit a plant response of 'susceptibility' (Box 1). Susceptibility is more likely to occur when the plant-herbivore association does not have an evolutionary history, and was the rationale for the 'new associations' model of agent selection (Hokkanen & Pimentel 1989). Susceptibility could also result if the weed has undergone evolutionary change post invasion resulting in a trade-off between growth and/or fecundity, and traits that enable resistance and tolerance (Blossey & Notzold 1995). The frequency of occurrence of such trade-offs is unclear, and it is often difficult to separate from plasticity in response to novel environments (Willis *et al.* 2000; Rogers & Siemann




Fig. 6. Elasticity analysis of the transition matrix for *Parkinsonia aculeata* in Central Australia. Each block in the diagram corresponds to a number in the transition matrix. The lighter the colour, the more sensitive the population growth is to a given proportional change in that number.

Box 1 A classification of plant response to herbivory and strategies of plants that confer resistance to minimise damage to tissues (adapted from Belsky *et al.* 1993) and definitions of terms (adapted from Strauss & Agrawal 1999).

Resistance – Plant trait that reduces the preference and/or performance of an herbivore.

Avoidance – A plant trait that reduces the likelihood of interacting with a herbivore or experiencing a negative effect.

Escape – A life-history trait that enables a plant to 'avoid' interactions with a herbivore (e.g. dispersal of seeds prior to build-up of seed predator populations (Raghu *et al.* 2005)).

Defence – A physical or chemical plant trait that enables the plant to deter herbivore damage and/or negatively influences the fitness of the herbivore (e.g. trichomes or presence of alkaloids).

Tolerance – The degree to which plant fitness is affected by herbivore damage, relative to an undamaged state.

Regrowth – The ability of a plant (through either physiological traits or storage organs) to recover from herbivore damage. Regrowth may result in a damaged plant being less fit (undercompensation) as fit (equal compensation) or more fit (overcompensation) in comparison with an undamaged plant.

Susceptibility – The inability of a plant to resist herbivore damage and thereby suffering strongly negative fitness consequences (e.g. reduced growth rate (individual and demographic), low reproductive output).

2002, 2003, 2004; Siemann & Rogers 2003; Bossdorf *et al.* 2004; Maron *et al.* 2004).

Exploration for weed biological control agents typically does not follow the new association model and exploration efforts are often restricted to the target plant's centre of origin (Goolsby *et al.* 2006), where potential agents are likely to have a coevolved interaction with the target plant. Therefore, plant response in these circumstances is likely to encompass the whole gamut of strategies outlined above, with different responses to different types of herbivory (Box 1). In such circumstances it is important to quantify plant response to herbivory to identify the type and extent of damage that an agent needs to inflict to achieve the desired management outcome (Raghu & Dhileepan 2005).

Is it possible to ascertain plant response to different types of herbivory *a priori*?

Once a vulnerable life stage of a weed has been identified, studying response to herbivory can yield valuable information in the context of agent selection. But is it possible to make these decisions without testing the response to every potential insect/pathogen identified on it in the native range? This can be a particularly daunting task in the case of biological control of weeds that have a diverse natural enemy complex. We propose that an alternate approach that simulates herbivory can provide valuable information for agent selection.

Simulated herbivory (mechanical damage to plants that approximates removal of tissues by a herbivore) has a long history of use in ecological research and has been a topic of vibrant debate (Baldwin 1990; Walling 2000; Hjalten 2004; Lehtilä & Boalt 2004). In a recent meta-analysis of 46 published papers on simulated herbivory, Lehtilä and Boalt (2004) found that 68% of the experiments showed differences in some measure of plant response between simulated and actual herbivory while 35% of the statistical analyses in those experiments showed a difference between simulated and actual herbivory. In studies that compared simulated and insect herbivory, 15% of the analyses revealed differences in plant response to the two methods of inflicting damage. When the types of plant response measured was examined, measures of growth and reproduction were less sensitive than other measures to differences in the damage methods with 26% of the analyses showing a difference between the two types of herbivory. Despite these differences, in only 3% of the studies did artificial and natural damage produce opposite plant responses, indicating that the direction of plant response to herbivory can be meaningfully inferred from simulated herbivory trials, while we need to be more cautious about interpreting the magnitude of plant response. In synthesising their findings, Lehtilä and Boalt (2004) concluded, 'the most important effects that artificial damage should succeed in simulating correctly are often the ones affecting biomass and population dynamics of interacting species'.

The advantages of simulated herbivory experiments include the ability to control for type, intensity and frequency of the damage, while simultaneously controlling other con-

foundering effects (Hjalten 2004). The obvious shortcomings include that it is difficult to precisely simulate natural herbivory in terms of the type, distribution and timing of damage and the interactions of damage with other important factors (Hjalten 2004; Lehtilä & Boalt 2004). Furthermore, certain types of physiological associations of herbivores (e.g. galling, other endophagous insects) cannot be simulated. This suggests that the value of simulated herbivory experiments rests with the type of questions being asked. While simulated herbivory may provide valuable clues about plant compensation or tolerance of damage, inferences on more complex ecological interactions are likely to be flawed (Hjalten 2004). Simulated herbivory experiments are less appropriate for understanding the influence of herbivory on higher trophic levels (e.g. parasitoid visitation in response to different plant injury levels), induced plant responses (where insect saliva may be important (Walling 2000)), effects of herbivory on soil processes or changes in competitive effects of plants suffering herbivory with those that are undamaged (Hjalten 2004; Lehtilä & Boalt 2004).

Studying plant response to simulated herbivory to different parts/modules can yield significant ecological insights about compensation and tolerance towards herbivory (Welter 1991; Marquis 1992; Hjalten *et al.* 1993; Welter & Steggall 1993; Foggo 1996; McLaren 1996; Gavloski & Lamb 2000; Tiffin & Inouye 2000; Rogers & Siemann 2002; Sullivan 2003; Hjalten 2004; Lehtilä & Boalt 2004; van Kleunen *et al.* 2004); information that could prove extremely useful in agent selection for weed biological control. Such an approach could help identify the guilds of herbivores most likely to have an impact on plant productivity and growth rates, thereby narrowing the prospective list of biological control agents, and prioritisation of agents in relation to management expectations (Ehler 1998). These studies could also provide guidance to exploration efforts, thereby allowing the limited time available for exploration in the native range to be targeted at the guilds most likely to yield effective agents. These results could also be integrated into models of weed-herbivore interactions to explore likely population level consequences of different types and levels of damage. At the very least, such studies generate hypotheses of the potential efficacy of the different guilds of herbivores (providing an explicit *a priori* basis for selection of particular agents) that can be tested against the performance of biological control agents during quarantine testing and after approval for release (Winder & van Emden 1980; Welter 1991; Broughton 2003; Sullivan 2003; Balciunas 2004; van Kleunen *et al.* 2004; Rebek & O'Neil 2005).

Several recent biological control projects in Australia have benefited from this approach. Simulated herbivory studies on *Mimosa pigra* (giant sensitive tree) in the Northern Territory have been used to generate *a priori* predictions for the performance and likely efficacy of *Macaria pallidata* (Wirt 2006a,b). Similar research on *Alternanthera phylloxeroides* (alligator weed) revealed that the effects of the beetle *Agasicles hygrophilla* could have been reliably predicted (Schooler *et al.* 2006). The response of *Jatropha gossypifolia* (bellyache bush) has provided valuable insights into the way agent

prioritisation decisions need to be made for this species (S Raghu unpubl. data 2005). Detailed experiments on *Macfadyena unguis-cati* (cat's claw creeper) have indicated that a defoliator that is likely to remove at least 75% of the leaf material twice over an 18-week period is the one most likely to control this liana. More significantly though, these simulated herbivory trials suggested that below-ground herbivory is likely to prove counter-productive to the objectives of biological control by triggering vigorous plant compensation. Based on this, leaf feeders are currently being screened for their safety and other guilds of herbivores have been ranked lower (Raghu & Dhileepan 2005; Raghu *et al.* 2006). Studies on *Parthenium hysterophorus* have confirmed that plant response to simulated and actual defoliation is comparable and, therefore, simulated herbivory could have been a tool used in the prioritisation of agents like *Zygogramma bicolorata* (Raghu & Dhileepan 2005). An important point to note is that all the above examples are for leaf-feeding herbivores, highlighting the earlier caveat that the simulated herbivory approach may be more appropriate for ectophagous than endophagous herbivores.

VERIFYING AGENT EFFICACY (FILTER 3)

Once clues about the life stage of the plant to target (Filter 1) and guild a successful herbivore is likely to come from (Filter 2) have been gathered, it is important to assess whether any among the pool of available herbivores can cause the desirable level of damage to achieve the management objective (i.e. our definition of 'successful management') (Fig. 1). There are two components to this. First, it is important to verify whether herbivores in the guild identified can cause the desirable level of damage (for examples of such research on temperate weeds, see Sheppard *et al.* 1994; Brieese 1996, 2000, 2006; Swirepik *et al.* 1996; Edwards & Brown 1997; Rees & Paynter 1997; Brieese & Zapater 2002; Brieese *et al.* 2002a,b, 2003; Zapater *et al.* 2004) and whether plant response to actual herbivory is similar to that identified by simulated herbivory studies (Hjalten 2004; Lehtilä & Boalt 2004). Second, to cause the desired level of damage, the candidate agent may need to attain a certain population density. The former is dealt with in this section, while the latter is dealt with in a subsequent paper (see Zalucki & van Klinken 2006).

At the early stages of a biocontrol program when agent selection decisions are made, any experiments using the candidate agents in the introduced range have to be conducted under quarantine conditions. Detailed tests on host range of agents are undertaken under the confines of quarantine. However, the data collection often stops with identifying host selection decisions made by the insect, and any effects on the plant are usually described qualitatively (e.g. Day *et al.* 1999; Dhileepan *et al.* 2005; Heard *et al.* 2005; Scott & Yeoh 2005). With a little more effort, it is possible to adapt such host-testing studies to gather quantitative data on level of herbivory an agent is able to inflict and the subsequent impacts on the plant in terms of regrowth potential, and in certain circumstances

reproductive output (S Raghu *et al.* unpubl. data 2005). Admittedly, the confines of quarantine and small cage testing are likely to overestimate or underestimate the effects of an agent if plants are healthy and growing in optimum conditions. Therefore, exclosure studies or observational data in the weed's native range during exploration efforts could yield additional insights to validate impacts of particular guilds of agents (Sheppard *et al.* 1994; Brieese 1996; Swirepik *et al.* 1996; Edwards & Brown 1997; Rees & Paynter 1997; Brieese 2000, 2006; Brieese & Zapater 2002; Brieese *et al.* 2002a,b, 2003; Zapater *et al.* 2004; Hafliger *et al.* 2005). Information on impacts can subsequently be fed-back into matrix models on plant demography to see how these affect population dynamics of the target weed in relation to our management objective. The integration of agent impacts into plant demography models have been used to assess non-target effects (e.g. Louda *et al.* 2005) and for agent evaluation studies (e.g. Shea & Kelly 1998), but can easily be adapted for agent selection.

CONCLUSION – PLACING THE FILTER OF EFFICACY BEFORE THE FILTER OF SAFETY

A central assumption in weed biological control is that specialist herbivores (typically insects, mites and pathogens) can exert a strong top-down influence on plant populations (McFadyen 1998, 2003) in the native range, and the escape from this influence during the process of the plant invading new regions is thought to lead to its invasiveness (referred to as the 'enemy release hypothesis'; Keane & Crawley 2002). This idea is seldom tested prospectively (Maron & Vila 2001) and is often validated retrospectively by the loss of herbivores and pathogens (Mitchell & Power 2003; Torchin *et al.* 2003) and through reference to successful biological control projects (McFadyen 1998, 2003). Through a combination of studies on plant demography and plant response to simulated and/or actual herbivory outlined above, the role of herbivores in regulating plant populations can be explicitly examined *a priori* and any potential role as biological control agents can be evaluated.

We suggest the following sequence of steps, based on plant demography and plant response to herbivory, for refining agent selection decisions in weed biological control.

- 1 Identify the life-history stage or transitions that when affected have the greatest negative consequence for plant demography.
- 2 For this vulnerable life stage, understand plant response to herbivory at the level of the individual plant:
 - Use approaches such as simulated herbivory or actual herbivory experiments in the native and introduced ranges to assess plant response (being aware that in the introduced range, the agent is unlikely to have specialist natural enemies).
 - Identify the type and magnitude of damage to which the plant is most vulnerable (i.e. produces the desired effects in relation to the management objective).

- 3 Prioritise an agent from the guild that causes the type of damage required in (2) to the life stage/transition identified in (1).
 - If (2) was done exclusively using simulated herbivory approaches, treat the findings as testable hypotheses and verify whether plant response to simulated damage and actual herbivore damage is comparable after importing the agent into quarantine, and if possible through native range studies.
- 4 Proceed to experimentally assessing the specificity of the agent.

We have adopted the metaphor of 'filters' to suggest how agent prioritisation may be improved. Access and taxonomic impediments may be more of a constraint on agent selection decisions in the case of neotropical weeds (see Goolsby *et al.* 2006) than temperate weeds. Therefore, simulating herbivory may be more of a tool in the context of selecting agents for the former. We are also intimately aware that in practice, logistic constraints may force scientists to be unable to subject candidate agents to these filters sequentially. This may not be as critical as ensuring that plant demography and plant response to herbivory be explored in significant empirical detail as a part of deciding which agents to release. As has been demonstrated in the case of research on temperate weeds (Sheppard *et al.* 1994; Briese 1996, 2000, 2006; Swirepik *et al.* 1996; Edwards & Brown 1997; Rees & Paynter 1997; Briese & Zapater 2002; Briese *et al.* 2002a,b, 2003; Zapater *et al.* 2004; Hafliger *et al.* 2005), such an approach offers the possibility of identifying agents that are likely to be effective and safe. At the very least such an approach makes the decisions behind the release of agents explicit, and these can be tested as hypotheses based on the release of agents.

ACKNOWLEDGEMENTS

David Briese, Harriet Hinz, Adele Reid and Rieks van Klinken made several detailed and insightful comments that challenged us to think very carefully about the ideas presented in this paper, and helped improve the paper significantly. We also thank Rieks van Klinken, Linda Anderson and Shane Campbell for providing unpublished data sets on *P. aculeata*.

REFERENCES

- Balciunas JK. 2004. Are mono-specific agents necessarily safe? The need for pre-release assessment of probable impact of candidate biocontrol agents, with some examples. In: *Proceedings of the XI International Symposium on Biological Control of Weeds* (eds JM Cullen, DT Briese, DJ Kriticos, WM Lonsdale, L Morin & JK Scott), pp. 252–257. CSIRO Entomology, Canberra, Australia.
- Baldwin IT. 1990. Herbivory simulations in ecological research. *Trends in Ecology and Evolution* **5**, 91–93.
- Belsky AJ, Carson WP, Jensen C & Fox JA. 1993. Overcompensation by plants: herbivore optimization or red herring? *Evolutionary Ecology* **7**, 109–121.
- Blossey B & Notzold R. 1995. Evolution of increased competitive ability in invasive nonindigenous plants: a hypothesis. *Journal of Ecology* **83**, 887–889.
- Bosssdorf O, Prati D, Auge H & Schmid B. 2004. Reduced competitive ability in an invasive plant. *Ecology Letters* **7**, 346–353.
- Briese DT. 1996. Potential impact of the stem-boring weevil *Lixus cardui* on the growth and reproductive capacity of *Onopordum* spp. thistles. *Biocontrol Science and Technology* **6**, 251–261.
- Briese DT. 2000. Impact of the *Onopordum* capitulum weevil *Larinus latus* on seed production by its host plant. *Journal of Applied Ecology* **37**, 238–246.
- Briese DT. 2004. Weed biological control: applying science to solve seemingly intractable problems. *Australian Journal of Entomology* **43**, 304–317.
- Briese DT. 2006. Can an *a priori* strategy be developed for biological control? The case of *Onopordum* spp thistles in Australia. *Australian Journal of Entomology* **45**, 317–323.
- Briese DT & Zapater M. 2002. A strategy for the biological control of blue heliotrope (*Heliotropium amplexicaule*). In: *Proceedings of the 13th Australian Weeds Conference* (eds H Spafford Jacob, J Dodd & JH Moore), pp. 394–397. Plant Protection Society of Western Australia, Perth, Australia.
- Briese DT, Petit W, Swirepik A & Walker A. 2002a. A strategy for the biological control of *Onopordum* spp. thistles in south-eastern Australia. *Biocontrol Science and Technology* **12**, 121–137.
- Briese DT, Thomann T & Vitou J. 2002b. Studies on the ecology of crown weevil, *Trichosirocalus briesei*, and its impact on growth and reproduction of its host *Onopordum* spp. thistles. *Journal of Applied Ecology* **39**, 688–698.
- Briese DT, Vitou J, Boettcher S & Thomann T. 2003. Prevalence and impact of the crown fly, *Botanophila spinosa*, on its host thistle, *Onopordum acanthium* in southern France. *Acta Oecologica* **24**, 77–86.
- Broughton S. 2003. Effect of artificial defoliation on growth and biomass of *Lantana camara* L. (Verbenaceae). *Plant Protection Quarterly* **18**, 110–115.
- Buckley YM, Rees M, Sheppard AW & Smyth MJ. 2005. Stable coexistence of an invasive plant and biocontrol agent: a parameterized coupled plant-herbivore model. *Journal of Applied Ecology* **42**, 70–79.
- Caswell H. 2001. *Matrix Population Models: Construction, Analysis and Interpretation*, 2nd edn. Sinauer Associates Inc., Sunderland, Massachusetts, USA.
- Crawley MJ, ed. 1997. *Plant Ecology*, 2nd edn. Blackwell Science, Cambridge, Massachusetts, USA.
- Cullen JH. 1992. Prediction and practice in biological control. In: *Proceedings of the First International Weed Control Congress*, Vol. 2 (ed. RG Richardson), pp. 147–150. Monash University, Melbourne, Australia.
- Cullen JM. 1995. Predicting effectiveness: fact and fantasy. In: *Proceedings of the VIII International Symposium on Biological Control of Weeds* (eds ES Delfosse & RR Scott), pp. 103–109. Agriculture Canada, Ottawa, Canada.
- Day MD, Willson BW & Nahrung HF. 1999. The life history and host range of the golden lantana beetle, *Charidotis pygmaea* (Col. Chrysomelidae), a biological control agent for *Lantana montevidensis* (Verbenaceae). *Biocontrol Science and Technology* **9**, 347–354.
- Denoth M, Frid L & Myers JH. 2002. Multiple agents in biological control: improving the odds? *Biological Control* **24**, 20–30.
- Dhileepan K, Treviño M, Donnelly GP & Raghu S. 2005. Risk to non-target plants from *Charidotis auroguttata* (Chrysomelidae: Coleoptera), a potential biocontrol agent for cat's claw creeper *Macfadyena unguis-cati* (Bignoniaceae) in Australia. *Biological Control* **32**, 450–460.
- Edwards PB & Brown EM. 1997. *Mesoclanis* seed flies (Diptera: Tephritidae) on *Chrysanthemoides* (Asteraceae) in South Africa: distribution, attack strategy and impact on seed production. *Bulletin of Entomological Research* **87**, 127–135.
- Ehler LE. 1998. Invasion biology and biological control. *Biological Control* **13**, 127–133.
- Foggo A. 1996. Long and short-term changes in plant growth following simulated herbivory: adaptive responses to damage? *Ecological Entomology* **21**, 198–202.
- Gavloski JE & Lamb RJ. 2000. Compensation by cruciferous plants is specific to the type of simulated herbivory. *Environmental Entomology* **29**, 1273–1282.

- Goolsby JA, van Klinken RD & Palmer WA. 2006. Maximising the contribution of native-range studies towards the identification and prioritisation of weed biocontrol agents. *Australian Journal of Entomology* **45**, 276–286.
- Hafliger P, Schwarzlaender M & Blossey B. 2005. Biology of *Platycephala planifrons* (Diptera: Chloropidae) and its potential effectiveness as biological control agent for invasive *Phragmites australis* in North America. *Biological Control* **34**, 302–311.
- Heard T, Paynter Q, Chan RR & Mira A. 2005. *Malacorhinus irregularis* for biological control of *Mimosa pigra*: host-specificity, life cycle, and establishment in Australia. *Biological Control* **32**, 252–262.
- Hjalten J. 2004. Simulating herbivory: problems and possibilities. In: *Insects and Ecosystem Function* (eds WW Weisser & E Siemann), pp. 243–256. Springer-Verlag, Berlin, Germany.
- Hjalten J, Danell K & Ericson L. 1993. Effects of simulated herbivory and intraspecific competition on the compensatory ability of birches. *Ecology* **74**, 1136–1142.
- Hoddle MS. 2004a. Restoring balance: using exotic species to control invasive exotic species. *Conservation Biology* **18**, 38–49.
- Hoddle MS. 2004b. The strength of biological control in the battle against invasive pests: a reply. *Conservation Biology* **18**, 61–64.
- Hokkanen H & Pimentel D. 1989. New associations in biological control: theory and practice. *Canadian Entomologist* **121**, 829–840.
- Howarth FG. 1991. Environmental implications of classical biological control. *Annual Review of Entomology* **36**, 485–509.
- Keane RM & Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology and Evolution* **17**, 164–170.
- Lehtilä K & Boalt E. 2004. The use and usefulness of artificial herbivory in plant-herbivore studies. In: *Insects and Ecosystem Function* (eds WW Weisser & E Siemann), pp. 257–275. Springer-Verlag, Berlin, Germany.
- Louda SM & Stiling P. 2004. The double-edged sword of biological control in conservation and restoration. *Conservation Biology* **18**, 50–53.
- Louda SM, Arnett AE, Rand TA & Russell FL. 2003a. Invasiveness of some biological control insects and adequacy of their ecological risk assessment and regulation. *Conservation Biology* **17**, 73–82.
- Louda SM, Pemberton RW, Johnson MT & Follett PA. 2003b. Nontarget effects – the Achilles' heel of biological control? Retrospective analyses to reduce risk associated with biocontrol introductions. *Annual Review of Entomology* **48**, 365–396.
- Louda SM, Rand TA, Arnett AE, McClay A, Shea K & McEachern AK. 2005. Evaluation of ecological risk to populations of a threatened plant from an invasive biocontrol insect. *Ecological Applications* **15**, 234–249.
- Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M & Bazzaz FA. 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* **10**, 689–710.
- Maron JL & Vila M. 2001. When do herbivores affect plant invasion? Evidence for the natural enemies and biotic resistance hypotheses. *Oikos* **95**, 361–373.
- Maron JL, Vila M & Arnason J. 2004. Loss of enemy resistance among introduced populations of St. John's Wort (*Hypericum perforatum*). *Ecology* **85**, 3243–3253.
- Marquis RJ. 1992. The selective impact of herbivores. In: *Plant Resistance to Herbivores and Pathogens: Ecology, Evolution and Genetics* (eds RS Fritz & EL Simms), pp. 301–325. University of Chicago Press, Chicago, Illinois, USA.
- Maywald GF, Sutherst RW & Zalucki MP. 1999. *DYMEX Professional: Modelling Natural Systems*. CSIRO, Melbourne, Australia.
- McClay A & Balciunas JK. 2006. The role of pre-release efficacy assessment in selecting classical biological control agents for weeds – applying the Anna Karenina principle. *Biological Control* **35**, 197–207.
- McEvoy PB. 2002. Insect–plant interactions on a planet of weeds. *Entomologia Experimentalis et Applicata* **104**, 165–179.
- McEvoy PB & Coombs EM. 1999. Biological control of plant invaders: regional patterns, field experiments, and structured population models. *Ecological Applications* **9**, 387–401.
- McEvoy PB & Coombs EM. 2000. Why things bite back: unintended consequences of biological control of weeds. In: *Non-target Effects of Biological Control* (eds PA Follett & JJ Duan), pp. 167–195. Kluwer Academic Publishers, Dordrecht, Netherlands.
- McFadyen REC. 1998. Biological control of weeds. *Annual Review of Entomology* **43**, 369–393.
- McFadyen RE. 2003. Does ecology help in the selection of biocontrol agents? In: *Improving the Selection, Testing and Evaluation of Weed Biological Control Agents* (eds H Spafford Jacob & DT Briese), pp. 5–9. Cooperative Research Centre for Australian Weed Management, Adelaide, Australia.
- McLaren B. 1996. Plant-specific response to herbivory: simulated browsing of suppressed balsam fir on Isle Royale. *Ecology* **77**, 228–235.
- Mitchell CE & Power AG. 2003. Release of invasive plants from fungal and viral pathogens. *Nature* **421**, 625–627.
- Pimentel D, ed. 2002. *Biological Invasions: Economic and Environmental Costs of Alien Plant, Animal, and Microbe Species*. CRC Press, Boca Raton, Florida, USA.
- Pimentel D, McNair S, Janecka J *et al.* 2001. Economic and environmental threats of alien plant, animal, and microbe invasions. *Agriculture, Ecosystems and Environment* **84**, 1–20.
- Raghu S & Dhileepan K. 2005. The value of simulating herbivory in selecting effective weed biological control agents. *Biological Control* **34**, 265–273.
- Raghu S, Wiltshire C & Dhileepan K. 2005. Predispersal seed predation in the invasive legume *Leucaena leucocephala* is limited by the duration of pod retention. *Austral Ecology* **30**, 310–318.
- Raghu S, Dhileepan K & Treviño M. 2006. Response of an invasive liana to simulated herbivory: implications for its biological control. *Acta Oecologica* **29**, 335–345.
- Rebek KA & O'Neil RJ. 2005. Impact of simulated herbivory on *Alliaria petiolata* survival, growth, and reproduction. *Biological Control* **34**, 284–289.
- Rees M & Paynter Q. 1997. Biological control of broom: modelling the determinants of abundance and potential impact of introduced insect herbivores. *Journal of Applied Ecology* **34**, 1203–1222.
- Rogers WE & Siemann E. 2002. Effects of simulated herbivory and resource availability on native and invasive exotic tree seedlings. *Basic and Applied Ecology* **3**, 297–307.
- Rogers WE & Siemann E. 2003. Effects of simulated herbivory and resources on Chinese tallow tree (*Sapium sebiferum*, Euphorbiaceae) invasion of native coastal prairie. *American Journal of Botany* **90**, 243–249.
- Rogers WE & Siemann E. 2004. Invasive ecotypes tolerate herbivory more effectively than native ecotypes of the Chinese tallow tree *Sapium sebiferum*. *Journal of Applied Ecology* **41**, 561–570.
- Schooler S, Baron Z & Julien M. 2006. Effect of simulated and actual herbivory on alligator weed, *Alternanthera philoxeroides*, growth and reproduction. *Biological Control* **36**, 74–79.
- Schroeder D & Goeden RD. 1986. The search for arthropod natural enemies of introduced weeds for biological control – theory and practice. *Biocontrol News and Information* **7**, 147–155.
- Scott JK & Yeoh PB. 2005. Biology and host specificity of *Apion minutum* (Coleoptera: Apionidae) from Israel, a potential biological control agent for *Emex australis* and *Emex spinosa* (Polygonaceae) in Australia. *Biological Control* **33**, 20–31.
- Shea K & Kelly D. 1998. Estimating biocontrol agent impact with matrix models: *Carduus nutans* in New Zealand. *Ecological Applications* **8**, 824–832.
- Shea K, Kelly D, Sheppard AW & Woodburn TL. 2005. Context dependent biological control of an invasive thistle. *Ecology* **86**, 3174–3181.
- Sheppard AW. 2003. Prioritising agents based on predicted efficacy: beyond the lottery approach. In: *Improving the Selection, Testing and Evaluation of Weed Biological Control Agents* (eds H Spafford Jacob & DT Briese), pp. 11–21. Cooperative Research Centre for Australian Weed Management, Adelaide, Australia.
- Sheppard AW & Raghu S. 2005. Working at the interface of art and science: how best to select an agent for classical biological control? *Biological Control* **34**, 233–235.
- Sheppard AW, Cullen JM & Aeschlimann J-P. 1994. Predispersal seed predation on *Carduus nutans* (Asteraceae) in southern Europe. *Acta Oecologica* **15**, 529–541.
- Sheppard AW, Hill R, DeClerck-Floate RA *et al.* 2003. A global review of risk-benefit-cost analysis for the introduction of classical

- biological control agents against weeds: a crisis in the making? *Biocontrol News and Information* **24**, 91N–108N.
- Siemann E & Rogers WE. 2003. Herbivory, disease, recruitment limitation and success of alien and native tree species. *Ecology* **84**, 1489–1505.
- Silvertown JW & Lovett Doust J. 1993. *Introduction to Plant Population Biology*. Blackwell Scientific Publications, Oxford, UK.
- Simberloff D & Stiling P. 1996. How risky is biological control? *Ecology* **77**, 1965–1974.
- Strauss SY & Agrawal AA. 1999. The ecology and evolution of plant tolerance to herbivory. *Trends in Ecology and Evolution* **14**, 179–185.
- Sullivan JJ. 2003. Density-dependent shoot-borer herbivory increases the age of first reproduction and mortality of neotropical tree saplings. *Oecologia* **136**, 96–106.
- Swirepik AE, Sheppard AW & Smyth MJ. 1996. *Meligethes planiusculus* (Nitidulidae): an inflorescence-feeding beetle of good potential for the biological control of Paterson's curse. In: *Proceedings of the 11th Australian Weeds Conference* (ed. RCH Shepherd), pp. 294–297. Weed Science Society of Victoria, Melbourne, Australia.
- Thomas MB & Willis AJ. 1998. Biocontrol – risky but necessary? *Trends in Ecology and Evolution* **13**, 325–329.
- Tiffin P & Inouye BD. 2000. Measuring tolerances to herbivory: accuracy and precision of estimates made using natural versus imposed damage. *Evolution* **54**, 1024–1029.
- Torchin ME, Lafferty KD, Dobson AP, McKenzie VJ & Kuris AM. 2003. Introduced species and their missing parasites. *Nature* **421**, 628–630.
- van Kleunen M, Ramponi G & Schmid B. 2004. Effects of herbivory simulated by clipping and jasmonic acid on *Solidago canadensis*. *Basic and Applied Ecology* **5**, 173–181.
- van Klinken RD. 2005. Total annual seed loss on a perennial legume through predation by insects: the importance of within-season seed and seed feeder dynamics. *Austral Ecology* **30**, 414–425.
- van Klinken RD & Raghu S. 2006. A scientific approach to agent selection. *Australian Journal of Entomology* **45**, 253–258.
- Walling LL. 2000. The myriad of plant responses to herbivores. *Journal of Plant Growth Regulation* **19**, 195–216.
- Welter SC. 1991. Responses of tomato to simulated and real herbivory by tobacco hornworm (Lepidoptera: Sphingidae). *Environmental Entomology* **20**, 1537–1541.
- Welter SC & Steggall JW. 1993. Contrasting the tolerance of wild and domesticated tomatoes to herbivory: agroecological implications. *Ecological Applications* **3**, 271–278.
- Willis AJ, Memmott J & Forrester RI. 2000. Is there evidence for the post-invasion evolution of increased size among invasive plant species? *Ecology Letters* **3**, 275–283.
- Winder JA & van Emden HF. 1980. Selection of effective biological control agents from artificial defoliation/insect cage experiments. In: *Proceedings of the Fifth International Symposium on Biological Control of Weeds* (ed. ES Delfosse), pp. 415–439. CSIRO, Melbourne, Australia.
- Wirf LA. 2006a. The effect of manual defoliation and *Macaria pallidata* (Geometridae) herbivory on *Mimosa pigra*: implications for biological control. *Biological Control* **37**, 346–353.
- Wirf LA. 2006b. Using simulated herbivory to predict the efficacy of a biocontrol agent – the effect of manual defoliation and *Macaria pallidata* (Geometridae) herbivory on *Mimosa pigra* seedlings. *Australian Journal of Entomology* **45**, 324–326.
- Zalucki M & van Klinken RD. 2006. Predicting population dynamics of weed biological control agents: science or gazing into crystal balls? *Australian Journal of Entomology* **45**, 331–344.
- Zapater M, Bartoloni N, Perez-Camargo G & Briece DT. 2004. Natural impact of the flea-beetle, *Longitarsus* spp., on *Heliotropium amplexicaule* in Argentina and its potential for use as a biological control agent in Australia. In: *Proceedings of the XI International Symposium on Biological Control of Weeds* (eds JM Cullen, DT Briece, DJ Kriticos, WM Lonsdale, L Morin & JK Scott), pp. 208–214. CSIRO Entomology, Canberra, Australia.

Accepted for publication 17 July 2006.